

Medical Information Request: ZUSDURI™ (mitomycin) for intravesical solution and Sodium Bicarbonate

Thank you for your question regarding the use of sodium bicarbonate as a pre-medication for ZUSDURI.

ZUSDURI™ is indicated for the treatment of adult patients with recurrent low-grade intermediate-risk non-muscle invasive bladder cancer (LG-IR-NMIBC).

Background:

ENVISION is a Phase 3, single-arm, multinational study evaluating the efficacy and safety of ZUSDURI (UGN-102) as a primary chemoablative therapy in 240 patients with LG-IR-NMIBC across 56 sites in the United States and Europe.

Utilizing UroGen's proprietary sterile hydrogel technology, ZUSDURI is a hydrogel-based formulation designed to enable longer exposure of bladder tissue to mitomycin. Based on patient-reported visibility of gel in urine post-treatment, ZUSDURI has a median dwell time of 5 hours with reports up to 24 hours. The reverse thermal properties of ZUSDURI allow for local administration of mitomycin as a liquid under chilled conditions, with subsequent conversion to a semisolid gel depot following instillation into the bladder.

Response:

In the ENVISION Phase 3 study, patients received ZUSDURI, 75 mg mitomycin in 56 mL admixture, once-weekly intravesically for 6 weeks. The study did not require sodium bicarbonate or other pre-medications as part of the protocol.

ZUSDURI Safety Profile:

In Phase 3 ENVISION,

- The most common ($\geq 10\%$) adverse reactions, including laboratory abnormalities, that occurred in patients treated with ZUSDURI were increased creatinine, increased potassium, dysuria, decreased hemoglobin, increased aspartate aminotransferase, increased alanine aminotransferase, increased eosinophils, decreased lymphocytes, urinary tract infection, decreased neutrophils, and hematuria.
- Serious adverse reactions occurred in 12% of patients who received ZUSDURI, including urinary retention (0.8%) and urethral stenosis (0.4%). A fatal adverse reaction of cardiac failure occurred in 1 (0.4%) patient (0.4%) receiving ZUSDURI.
- Dosage interruption of ZUSDURI due to adverse reactions occurred in 10% of patients. Adverse reactions ($\geq 2\%$) which required dosage interruption were urinary tract infection (2.5%) and dysuria (2.5%)
- Permanent discontinuation of ZUSDURI due to an adverse reaction occurred in 2.9% of patients, including 1.7% who discontinued due to a renal or urinary disorder.

Please refer to the Full Prescribing Information for ZUSDURI [here](#).

ZUSDURI IMPORTANT SAFETY INFORMATION:

Contraindications

ZUSDURI is contraindicated in patients with perforation of the bladder or in patients with prior hypersensitivity reactions to mitomycin or any component of the product.

Warnings and Precautions

Risks in Patients with Perforated Bladder

ZUSDURI may lead to systemic exposure to mitomycin and severe adverse reactions if administered to patients with a perforated bladder or to those in whom the integrity of the bladder mucosa has been compromised. Evaluate the bladder before the intravesical instillation of ZUSDURI and do not administer to patients with a perforated bladder or mucosal compromise until bladder integrity has been restored.

Embryo-Fetal Toxicity

Based on findings in animals and mechanism of action, ZUSDURI can cause fetal harm when administered to a

pregnant woman. In animal reproduction studies, administration of mitomycin resulted in teratogenicity. Advise females of reproductive potential to use effective contraception during treatment with ZUSDURI and for 6 months following the last dose. Advise male patients with female partners of reproductive potential to use effective contraception during treatment with ZUSDURI and for 3 months following the last dose.

Adverse Reactions

Common Adverse Reactions

The most common ($\geq 10\%$) adverse reactions, including laboratory abnormalities, that occurred in patients treated with ZUSDURI were increased creatinine, increased potassium, dysuria, decreased hemoglobin, increased aspartate aminotransferase, increased alanine aminotransferase, increased eosinophils, decreased lymphocytes, urinary tract infection, decreased neutrophils, and hematuria.

Additional Adverse Reactions Information

Clinically relevant adverse reactions occurring in $< 10\%$ of patients who received ZUSDURI included increased urinary frequency, fatigue, urinary incontinence, urinary retention, urethral stenosis, genital pain, urinary urgency, genital edema, genital pruritus, genital rash, urethritis, acute kidney injury, balanoposthitis, and nocturia.

Use in Specific Populations

Lactation

Because of the potential for serious adverse reactions in a breastfed child, advise women not to breastfeed during treatment with ZUSDURI and for 1 week following the last dose.

Preparation and Administration Information

ZUSDURI is to be administered by intravesical instillation only. Do not administer ZUSDURI by pyelocalyceal instillation or by any other route.

ZUSDURI must be prepared and administered by a healthcare provider. To ensure proper dosing, it is important to follow the preparation instructions found in the ZUSDURI Instructions for Pharmacy and administration instructions found in the ZUSDURI Instructions for Administration.

ZUSDURI may discolor urine to a violet to blue color following the instillation procedure. Advise patients for at least 24 hours post-instillation to avoid urine contact with skin, to void urine sitting on a toilet, and to flush the toilet several times after use. Advise patients to wash hands, perineum or glans with soap and water after each instillation procedure.

ZUSDURI is a hazardous drug. Follow applicable special handling and disposal procedures.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit <http://www.fda.gov/medwatch> or call 1-800-FDA-1088. You may also report side effects to UroGen Pharma at 1-855-987-6436.

Please see accompanying Full Prescribing Information, Instructions for Pharmacy and Instructions for Administration.

References:

1. ZUSDURI™ (mitomycin) for intravesical solution. Prescribing Information. UroGen Pharma; 2025.
2. ZUSDURI™ (mitomycin) for intravesical solution. Instructions for Pharmacy (IFP)
3. ZUSDURI™ (mitomycin) for intravesical solution. Instructions for Administration (IFA)



4. Prasad SM, Shishkov D, Mihaylov NV, et al. Primary chemoablation of recurrent low-grade intermediate-risk nonmuscle-invasive bladder cancer with UGN-102: A single-arm, open-label phase 3 trial (ENVISION). *J Urol.* 2025;213(2):205-16.
5. Data on file. UroGen Pharma.

ZUSDURI™ is a trademark and UroGen® is a registered trademark of UroGen Pharma, Ltd.